

Concluding talk: Fundamental lessons and challenges from neutrinos

G. ALTARELLI

*Dipartimento di Matematica e Fisica, Università di Roma Tre, INFN, Sezione di Roma Tre, Via della Vasca Navale 84, I-00146 Rome, Italy and
CERN, Department of Physics, Theory Unit - CH-1211 Geneva 23, Switzerland*

Summary. — We present a concise review of the experimental developments on neutrino mixing and their theoretical implications as presented and discussed at this Conference. The recent data disfavour many models but the surviving ones still span a wide range going from Anarchy (no structure, no symmetry in the neutrino sector) to a maximum of symmetry, as for the models based on discrete non-abelian flavour groups which can be improved following the indications from the recent data.

1. – Introduction

Bruno Pontecorvo has pioneered the physics of neutrinos in many different aspects, as it has been impressively reviewed at this Conference [1]. In the last two decades experiments have established the existence of neutrino oscillations that Bruno had anticipated and the most important related parameters have been measured. These results represent a major progress of great importance for particle physics and cosmology. Neutrino physics is at present a most vital domain of particle physics and cosmology (with implications also for geology [2], cosmic rays and astronomy [3]) and the existing open questions are of crucial importance. In this concluding talk I will review the main lessons that we have learnt so far from neutrinos and discuss the present challenges in this field.

The main facts from ν mass and mixing [4] are that ν 's are not all massless but their masses are very small; probably their masses are small because ν 's are Majorana fermions with masses inversely proportional to the large scale M of interactions that violate lepton number (L) conservation. From the see-saw formula [5] together with the observed atmospheric oscillation frequency and a Dirac mass m_D of the order of the Higgs VEV, it follows that the Majorana mass scale $M \sim m_{\nu R}$ is empirically close to $10^{14} - 10^{15}$ GeV $\sim M_{GUT}$, so that ν masses fit well in the Grand Unification Theory (GUT) picture. Decays of ν_R with CP and L violation can produce a sizable B-L asymmetry that survives instanton effects at the electroweak scale thus explaining baryogenesis as arising from leptogenesis. There is still no direct proof that neutrinos are Majorana fermions:

detecting neutrino-less double beta decay ($0\nu\beta\beta$) would prove that ν 's are Majorana particles and that L is violated. It also appears that the active ν 's are not a significant component of dark matter in the Universe.

On the experimental side the main recent developments on neutrino mixing [4] were the results on θ_{13} [6, 7] from T2K, MINOS, DOUBLE CHOOZ, RENO and especially DAYA-BAY. The different experiments are in good agreement and the most precise is DAYA-BAY with the result $\sin^2 2\theta_{13} = 0.090_{-0.009}^{+0.008}$ [6] (equivalent to $\sin^2 \theta_{13} \sim 0.023 \pm 0.002$ or $\theta_{13} \sim (8.7 \pm 0.6)^\circ$). A summary of recent global fits to the data on oscillation parameters is presented in Table 1 [8], [9], [10]. The combined value of $\sin^2 \theta_{13}$ is by now about 10σ away from zero and the central value is rather large, close to the previous upper bound. In turn a sizable θ_{13} allows to extract an estimate of θ_{23} from accelerator data like T2K and MINOS. There are now solid indications of a deviation of θ_{23} from the maximal value, probably in the first octant [8]. In addition, some tenuous hints that $\cos \delta_{CP} < 0$ are starting to appear in the data.

TABLE I. – *Fits to neutrino oscillation data. For $\sin^2 \theta_{23}$ from Ref. [9] only the absolute minimum in the first octant is shown*

Quantity	Ref. [8]	Ref. [9]
Δm_{sun}^2 (10^{-5} eV ²)	$7.54_{-0.22}^{+0.26}$	7.50 ± 0.185
Δm_{atm}^2 (10^{-3} eV ²)	$2.43_{-0.10}^{+0.06}$	$2.47_{-0.067}^{+0.069}$
$\sin^2 \theta_{12}$	$0.307_{-0.016}^{+0.018}$	0.30 ± 0.013
$\sin^2 \theta_{23}$	$0.386_{-0.021}^{+0.024}$	$0.41_{-0.025}^{+0.037}$
$\sin^2 \theta_{13}$	0.0241 ± 0.025	0.023 ± 0.0023

A hot issue is the possible existence of sterile neutrinos [11] (see sect. 6).

2. – Neutrino masses and lepton number violation

Neutrino oscillations imply non vanishing neutrino masses which in turn demand either the existence of right-handed (RH) neutrinos (Dirac masses) or lepton number L violation (Majorana masses) or both. Given that neutrino masses are extremely small, it is really difficult from the theory point of view to avoid the conclusion that L conservation must be violated. In fact, in terms of lepton number violation the smallness of neutrino masses can be explained as inversely proportional to the very large scale where L is violated, of order M_{GUT} or even M_{Pl} .

If L conservation is violated neutrinos can be Majorana fermions. For a Majorana neutrino each mass eigenstate with given helicity coincides with its own antiparticle with the same helicity. As well known, for a charged massive fermion there are four states differing by their charge and helicity (the four components of a Dirac spinor) as required by Lorentz and CPT invariance. For a massive Majorana neutrino, neutrinos and antineutrinos can be identified and only two components are needed to satisfy the Lorentz and CPT invariance constraints. Neutrinos can be Majorana fermions because, among the fundamental fermions (i.e. quarks and leptons), they are the only electrically neutral ones. If, and only if, the lepton number L is not conserved, i.e. it is not a good quantum number, then neutrinos and antineutrinos can be identified. For Majorana neutrinos both Dirac mass terms, that conserve L ($\nu \rightarrow \nu$), and Majorana mass terms,

that violate L by two units ($\nu \rightarrow \bar{\nu}$), are in principle possible. Of course the restrictions from gauge invariance must be respected. So, for neutrinos the Dirac mass terms ($\bar{\nu}_R \nu_L + \text{h.c.}$) arise from the couplings with the Higgs field, as for all quarks and leptons. For Majorana masses, a $\nu_L^T \nu_L$ mass term has weak isospin 1 and needs two Higgs fields to make an invariant. On the contrary a $\nu_R^T \nu_R$ mass term is a gauge singlet and needs no Higgs. As a consequence, the RH neutrino Majorana mass M_R is not bound to be of the order of the electroweak symmetry breaking (induced by the Higgs vacuum expectation value) and can be very large (see below).

Some notation: the charge conjugated of ν is ν^c , given by $\nu^c = C(\bar{\nu})^T$, where $C = i\gamma_2\gamma_0$ is the charge conjugation matrix acting on the spinor indices. In particular $(\nu^c)_L = C(\bar{\nu}_R)^T$, so that, instead of using ν_L and ν_R , we can refer to ν_L and $(\nu^c)_L$, or simply ν and ν^c .

Once we accept L non-conservation we gain an elegant explanation for the smallness of neutrino masses. If L is not conserved, even in the absence of heavy RH neutrinos, Majorana masses for neutrinos can be generated by dimension five operators of the form

$$(1) \quad O_5 = \frac{(Hl)_i^T \lambda_{ij} (Hl)_j}{\Lambda} \quad ,$$

with H being the ordinary Higgs doublet, l_i the SU(2) LH lepton doublets, λ a matrix in flavour space, Λ a large scale of mass, possibly of order M_{GUT} or M_{Pl} and a charge conjugation matrix C between the lepton fields is understood. Neutrino masses generated by O_5 are of the order $m_\nu \approx v^2/\Lambda$ for $\lambda_{ij} \approx O(1)$, where $v \sim O(100 \text{ GeV})$ is the vacuum expectation value of the ordinary Higgs.

We consider that the existence of RH neutrinos ν^c is quite plausible because most GUT groups larger than SU(5) require them. In particular the fact that ν^c completes the representation 16 of SO(10): $16 = \bar{5} + 10 + 1$, so that all fermions of each family are contained in a single representation of the unifying group, is too impressive not to be significant. At least as a classification group SO(10) must be of some relevance in a more fundamental layer of the theory! Thus in the following we assume both that ν^c exist and that L is not conserved. With these assumptions the see-saw mechanism [5] is possible. We recall, also to fix notations, that in its simplest form it arises as follows. Consider the SU(3) \times SU(2) \times U(1) invariant Lagrangian giving rise to Dirac and ν^c Majorana masses (for the time being we consider the ν (versus ν^c) Majorana mass terms as comparatively negligible):

$$(2) \quad \mathcal{L} = -\nu^{cT} y_\nu (Hl) + \frac{1}{2} \nu^{cT} M \nu^c + \text{h.c.}$$

The Dirac mass matrix $m_D \equiv y_\nu v/\sqrt{2}$, originating from electroweak symmetry breaking, is, in general, non-hermitian and non-symmetric, while the Majorana mass matrix M is symmetric, $M = M^T$. We expect the eigenvalues of M to be of order M_{GUT} or more because ν^c Majorana masses are SU(3) \times SU(2) \times U(1) invariant, hence unprotected and naturally of the order of the cutoff of the low-energy theory. Since all ν^c are very heavy we can integrate them away and the resulting neutrino mass matrix reads:

$$(3) \quad m_\nu = m_D^T M^{-1} m_D \quad .$$

This is the well known see-saw mechanism result [5]: the light neutrino masses are quadratic in the Dirac masses and inversely proportional to the large Majorana mass.

If some ν^c are massless or light they would not be integrated away but simply added to the light neutrinos. Note that for $m_\nu \approx \sqrt{\Delta m_{atm}^2} \approx 0.05$ eV (see Table(1)) and $m_\nu \approx m_D^2/M$ with $m_D \approx v \approx 200$ GeV we find $M \approx 10^{15}$ GeV which indeed is an impressive indication for M_{GUT} .

If additional non-renormalizable contributions to O_5 , eq. (1), are comparatively non-negligible, they should simply be added. For instance in SO(10) or in left-right extensions of the SM, an $SU(2)_L$ triplet can couple to two lepton doublets and to two Higgs and may induce a sizeable contribution to neutrino masses. At the level of the low-energy effective theory, such contribution is still described by the operator O_5 of eq. (1), obtained by integrating out the heavy $SU(2)_L$ triplet. This contribution is called type II to be distinguished from that obtained by the exchange of RH neutrinos (type I). One can also have the exchange of a fermionic $SU(2)_L$ triplet coupled to a lepton doublet and a Higgs (type III). After elimination of the heavy fields, at the level of the effective low-energy theory, the three types of see-saw terms are equivalent. In particular they have identical transformation properties under a chiral change of basis in flavour space. The difference is, however, that in type I see-saw mechanism, the Dirac matrix m_D is presumably related to ordinary fermion masses because they are both generated by the Higgs mechanism and both must obey GUT-induced constraints. Thus more constraints are implied if one assumes the see-saw mechanism in its simplest type I version.

3. – Basic formulae for three-neutrino mixing

In this section we assume that there are only two distinct neutrino oscillation frequencies, the atmospheric [13] and the solar frequencies [12]. These two can be reproduced with the known three light neutrino species (with no need of sterile neutrinos).

Neutrino oscillations are due to a misalignment between the flavour basis, $\nu' \equiv (\nu_e, \nu_\mu, \nu_\tau)$, where ν_e is the partner of the mass and flavour eigenstate e^- in a left-handed (LH) weak isospin $SU(2)$ doublet (similarly for ν_μ and ν_τ) and the mass eigenstates $\nu \equiv (\nu_1, \nu_2, \nu_3)$ [14-16]:

$$(4) \quad \nu' = U\nu \quad ,$$

where U is the unitary 3 by 3 mixing matrix. Given the definition of U and the transformation properties of the effective light neutrino mass matrix m_ν in eq. (1):

$$(5) \quad \begin{aligned} \nu'^T m_\nu \nu' &= \nu^T U^T m_\nu U \nu \\ U^T m_\nu U &= \text{Diag}(m_1, m_2, m_3) \equiv m_{diag} \quad , \end{aligned}$$

we obtain the general form of m_ν (i.e. of the light ν mass matrix in the basis where the charged lepton mass is a diagonal matrix):

$$(6) \quad m_\nu = U^* m_{diag} U^\dagger \quad .$$

The matrix U can be parameterized in terms of three mixing angles θ_{12} , θ_{23} and θ_{13} ($0 \leq \theta_{ij} \leq \pi/2$) and one phase φ ($0 \leq \varphi \leq 2\pi$) [17], exactly as for the quark mixing matrix V_{CKM} . The following definition of mixing angles can be adopted:

$$(7) \quad U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{i\varphi} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\varphi} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where $s_{ij} \equiv \sin \theta_{ij}$, $c_{ij} \equiv \cos \theta_{ij}$. In addition, if ν are Majorana particles, we have the relative phases among the Majorana masses m_1 , m_2 and m_3 . If we choose m_3 real and positive, these phases are carried by $m_{1,2} \equiv |m_{1,2}|e^{i\phi_{1,2}}$ [18]. Thus, in general, 9 parameters are added to the SM when non-vanishing neutrino masses are included: 3 eigenvalues, 3 mixing angles and 3 CP violating phases.

In our notation the two frequencies, $\Delta m_I^2/4E$ ($I=\text{sun,atm}$), are parametrized in terms of the ν mass eigenvalues by

$$(8) \quad \Delta m_{sun}^2 \equiv |\Delta m_{12}^2|, \quad \Delta m_{atm}^2 \equiv |\Delta m_{23}^2| \quad .$$

where $\Delta m_{12}^2 = |m_2|^2 - |m_1|^2 > 0$ (positive by the definition of $m_{1,2}$) and $\Delta m_{23}^2 = m_3^2 - |m_2|^2$. The numbering 1,2,3 corresponds to our definition of the frequencies and in principle may not coincide with the ordering from the lightest to the heaviest state. In fact, the sign of Δm_{23}^2 is not known [a positive (negative) sign corresponds to normal (inverse) hierarchy]. The determination of the hierarchy pattern together with the measurement of the CP violating phase φ are among the main experimental challenges for future accelerators [19].

Oscillation experiments do not provide information about the absolute neutrino mass scale. Limits on that are obtained [4] from the endpoint of the tritium beta decay spectrum, from cosmology and from neutrinoless double beta decay ($0\nu\beta\beta$). From tritium we have an absolute upper limit of 2.2 eV (at 95% C.L.) [20] on the antineutrino mass eigenvalues involved in beta decay, which, combined with the observed oscillation frequencies under the assumption of three CPT-invariant light neutrinos, also amounts to an upper bound on the masses of the other active neutrinos. The future of the tritium measurement is the KATRIN experiment whose goal is to improve the present limit by about an order of magnitude [20]. Complementary information on the sum of neutrino masses is also provided by cosmology [21]. For the sum of all (quasi) stable (thermalized) neutrino masses the Planck experiment, also using the WMAP 9 and BAO data, finds the limit $\sum m_\nu \leq 0.23$ at 95% c.l. [22]. The discovery of $0\nu\beta\beta$ decay would be very important, as discussed in the next section, and would also provide direct information on the absolute scale of neutrino masses [23].

4. – Importance of neutrino-less double beta decay

The detection of neutrino-less double beta decay [23] would provide direct evidence of L non conservation and of the Majorana nature of neutrinos. It would also offer a way to possibly disentangle the 3 cases of degenerate, normal or inverse hierarchy neutrino spectrum. The quantity which is bound by experiments on $0\nu\beta\beta$ is the 11 entry of the ν mass matrix, which in general, from $m_\nu = U^* m_{diag} U^\dagger$, is given by :

$$(9) \quad |m_{ee}| = |(1 - s_{13}^2) (m_1 c_{12}^2 + m_2 s_{12}^2) + m_3 e^{2i\phi} s_{13}^2|$$

where $m_{1,2}$ are complex masses (including Majorana phases) while m_3 can be taken as real and positive and ϕ is the U phase measurable from CP violation in oscillation experiments. Starting from this general formula it is simple to derive the bounds for degenerate, inverse hierarchy or normal hierarchy mass patterns.

At present the best limits from the searches with Ge lead to $|m_{ee}| \sim (0.25 - 0.98)$ eV (GERDA,+HM+IGEX) and with Xe to $|m_{ee}| \sim (0.12 - 0.25)$ eV (EXO+Kamland Zen), where ambiguities on the nuclear matrix elements lead to the ranges shown [23].

In the next few years, experiments (CUORE, GERDA II, SNO+....) will reach a larger sensitivity on $0\nu\beta\beta$ by about an order of magnitude. Assuming the standard mechanism through mediation of a light massive Majorana neutrino, if these experiments will observe a signal this would indicate that the inverse hierarchy is realized, if not, then the normal hierarchy case still would remain a possibility.

5. – Baryogenesis via leptogenesis from heavy ν^c decay

In the Universe we observe an apparent excess of baryons over antibaryons. It is appealing that one can explain the observed baryon asymmetry by dynamical evolution (baryogenesis) starting from an initial state of the Universe with zero baryon number. For baryogenesis one needs the three famous Sakharov conditions: B violation, CP violation and no thermal equilibrium. In the history of the Universe these necessary requirements have possibly occurred at different epochs. Note however that the asymmetry generated during one such epoch could be erased in following epochs if not protected by some dynamical reason. In principle these conditions could be fulfilled in the SM at the electroweak phase transition. In fact, when kT is of the order of a few TeV, B conservation is violated by instantons (but B-L is conserved), CP symmetry is violated by the CKM phase and sufficiently marked out-of-equilibrium conditions could be realized during the electroweak phase transition. So the conditions for baryogenesis at the weak scale in the SM superficially appear to be present. However, a more quantitative analysis [24] shows that baryogenesis is not possible in the SM because there is not enough CP violation and the phase transition is not sufficiently strong first order, because the Higgs mass is too heavy. In SUSY extensions of the SM, in particular in the MSSM, there are additional sources of CP violation but also this possibility has by now become at best marginal after the results from LEP2 and the LHC.

If baryogenesis at the weak scale is excluded by the data it can occur at or just below the GUT scale, after inflation. But only that part with $|B - L| > 0$ would survive and not be erased at the weak scale by instanton effects. Thus baryogenesis at $kT \sim 10^{10} - 10^{15}$ GeV needs B-L violation and this is also needed to allow m_ν if neutrinos are Majorana particles. The two effects could be related if baryogenesis arises from leptogenesis then converted into baryogenesis by instantons [25]. The decays of heavy Majorana neutrinos (the heavy eigenstates of the see-saw mechanism) happen with violation of lepton number L, hence also of B-L and can well involve a sufficient amount of CP violation. Recent results on neutrino masses are compatible with this elegant possibility. Thus the case of baryogenesis through leptogenesis has been boosted by the recent results on neutrinos.

6. – Sterile neutrinos?

A number of hints have been recently collected for the existence of sterile neutrinos [11], that is neutrinos with no weak interactions (for a review see ref. [27]). They do not make yet an evidence but certainly pose an experimental problem that needs clarification (see, for example, Ref. [28]).

The MiniBooNE experiment published [29] a combined analysis of ν_e appearance in a ν_μ beam together with $\bar{\nu}_e$ appearance in a $\bar{\nu}_\mu$ beam. They observe an excess of events from neutrinos over expected background in the low energy region (below 500 MeV) of the event spectrum. In the most recent data the shapes of the neutrino and anti-neutrino spectra appear to be consistent with each other, showing excess events below 500 MeV

and data consistent with background in the high energy region. The allowed region from MiniBooNE anti-neutrino data has some overlap with the parameter region preferred by LSND. Recently the ICARUS experiment at Gran Sasso has published the results of a search for electrons produced by the CERN neutrino beam [30]. No excess over the background was observed. As a consequence a large portion of the region allowed by LSND, MiniBooNE, KARMEN... is now excluded.

Then there are $\bar{\nu}_e$ disappearance experiments: in particular, the reactor and the gallium anomalies. A reevaluation of the reactor flux [31] produced an apparent gap between the theoretical expectations and the data taken at small distances from reactors (≤ 100 m). A different analysis confirmed the normalization shift [32]. Similarly the Gallium anomaly [33] depends on the assumed cross-section which could be questioned.

These data hint at one or more sterile neutrinos with mass around 1 eV which would represent a major discovery in particle physics. Cosmological data allow for one single sterile neutrino but more than one are disfavoured by the stringent bounds arising from nucleosynthesis (assuming fully thermalized sterile neutrinos) [26]. Actually the recently published Planck data [22] on the cosmic microwave background (CMB) are completely consistent with no sterile neutrinos (they quote $N_{eff} = 0.30 \pm 0.27$). The absence of a positive signal in ν_μ disappearance in accelerator experiments (CDHSW, MINOS, CCFR, MiniBooNE-SciBooNE) creates a tension with LSND (if no CP viol.). For example, in 3+1 models there is a tension between appearance (LSND, MiniBooNe.....) and disappearance (MINOS...) [34]. However, a better 3+1 fit is obtained if the low energy MiniBooNe data are not included [11, 35]. In 3+1 models the short baseline reactor data and the gallium anomaly are not in tension with the other measurements. Fits with 2 sterile neutrinos do not solve all the tensions [34, 36]. In general in all fits the resulting sterile neutrino masses are too large when compared with the cosmological bounds on the sum of neutrino masses, if the contribution of the sterile neutrinos to the effective number of relativistic degrees of freedom is close to one.

In conclusion, the situation is at present confuse but the experimental effort should be continued because establishing the existence of sterile neutrinos would be a great discovery (an experiment to clarify the issue of sterile neutrinos is proposed on the CERN site [37]). In fact a sterile neutrino is an exotic particle not predicted by the most popular models of new physics.

As only a small leakage from active to sterile neutrinos is allowed by present neutrino oscillation data (see, for example, refs. [38, 39, 34, 40] and references therein), in the following we restrict our discussion to 3-neutrino models.

7. – Models of neutrino mixing

A long list of models have been formulated over the years to understand neutrino masses and mixings. With the continuous improvement of the data most of the models have been discarded by experiment. But the surviving models still span a wide range going from a maximum of symmetry, with discrete non-abelian flavour groups, to the opposite extreme of anarchy.

The rather large measured value of θ_{13} , close to the old CHOOZ bound and to the Cabibbo angle, and the indication that θ_{23} is not maximal both go in the direction of models based on Anarchy [41, 42], i.e. the idea that perhaps no symmetry is needed in the neutrino sector, only chance (this possibility has been recently reiterated, for example, in Ref. [43]). Anarchy can be formulated in a $SU(5) \otimes U(1)_{FN}$ context by taking different Froggatt-Nielsen [44] charges only for the $SU(5)$ tenplets (for example $10 \sim (a, b, 0)$,

where $a > b > 0$ is the charge of the first generation, b of the second, zero of the third) while no charge differences appear in the $\bar{5}$ (e. g. $\bar{5} \sim (0, 0, 0)$). The observed fact that the up-quark mass hierarchies are more pronounced than for down-quark and charged leptons is in agreement with this assignment. In models with no see-saw, the $\bar{5}$ charges completely fix the hierarchies (or Anarchy, if the case) in the neutrino mass matrix. If RH neutrinos are added, they transform as $SU(5)$ singlets and can in principle carry $U(1)_{FN}$ charges, which also, in the Anarchy case, must be all equal. With RH neutrinos the see-saw mechanism can take place and the resulting phenomenology is modified. The embedding of Anarchy in the $SU(5) \otimes U(1)_{FN}$ context allows to implement a parallel treatment of quarks and leptons. Note that implementing Anarchy and its variants in $SO(10)$ is difficult.

The $SU(5)$ generators act vertically inside one generation, whereas the $U(1)_{FN}$ charges differ horizontally from one generation to the other. If, for a given interaction vertex, the $U(1)_{FN}$ charges do not add to zero, the vertex is forbidden in the symmetric limit. However, the $U(1)_{FN}$ symmetry (that one can assume to be a gauge symmetry) is spontaneously broken by the VEVs v_f of a number of flavon fields with non-vanishing charge and GUT-scale masses. Then a forbidden coupling is rescued but is suppressed by powers of the small parameters $\lambda = v_f/M$, with M a large mass, with the exponents larger for larger charge mismatch. Thus the charges fix the powers of λ , hence the degree of suppression of all elements of mass matrices, while arbitrary coefficients k_{ij} of order 1 in each entry of mass matrices are left unspecified (so that the number of order 1 parameters exceeds the number of observable quantities). A random selection of these k_{ij} parameters leads to distributions of resulting values for the measurable quantities. For Anarchy the mass matrices in the neutrino sector (determined by the $\bar{5}$ and 1 charges) are totally random, while in the presence of unequal charges different entries carry different powers of the order parameter and thus some hierarchies are enforced.

Within this framework there are many variants of these models: fermion charges can all be nonnegative with only negatively charged flavons, or there can be fermion charges of different signs with either flavons of both charges or only flavons of one charge. In Ref. [45], given the new experimental results, we have made a reappraisal of Anarchy and its variants within the $SU(5) \times U(1)_{FN}$ GUT framework. Based on the most recent data we argue that the Anarchy ansatz is probably oversimplified and, in any case, not compelling. In fact, suitable differences of $U(1)_{FN}$ charges, if also introduced within pentaplets and singlets, lead to distributions that are in better agreement with the data with the same number of random parameters as for Anarchy. The hierarchy of quark masses and mixing and of charged lepton masses in all cases impose a hierarchy-defining parameter of the order of $\lambda_C = \sin \theta_C$, with θ_C being the Cabibbo angle. The weak points of Anarchy are that all mixing angles should be of the same order, so that the relative smallness of $\theta_{13} \sim o(\lambda_C)$ is not automatic. Similarly the smallness of $r = \Delta m_{solar}^2 / \Delta m_{atm}^2$ is not easily reproduced: with no See-Saw r is of $o(1)$, while in the See-Saw version of Anarchy the problem is only partially alleviated by the spreading of the neutrino mass distributions that follows from the product of three matrix factors in the See-Saw formula. An advantage is already obtained if Anarchy is only restricted to the 23 sector of leptons. In this case, with or without See-Saw, θ_{13} is naturally suppressed and, with a single fine tuning one gets both θ_{12} large and r small (this model was also recently rediscussed in Ref. [46]). Actually in Ref. [45] we have shown, for example, that the freedom of adopting RH neutrino charges of both signs, can be used to obtain a completely natural model where all small quantities are suppressed by the appropriate power of λ_C . In this model a lopsided Dirac mass matrix is combined with a generic Majorana matrix to

produce a neutrino mass matrix where the 23 subdeterminant is suppressed and thus r is naturally small and θ_{23} is large. In addition also θ_{12} is large while θ_{13} is suppressed. We stress again that the number of random parameters is the same in all these models: one coefficient of $o(1)$ for every matrix element. Moreover, with an appropriate choice of charges, it is not only possible to reproduce the charged fermion hierarchies and the quark mixing, but also the order of magnitude of all small observed parameters can be naturally guaranteed. In conclusion, models based on chance are still perfectly viable, but we consider Anarchy a particularly simple choice perhaps oversimplified and certainly not compelling and we have argued that the hierarchy of charged fermion masses needs a minimum of flavour symmetry (like $U(1)_{FN}$) which, to some extent, can also be effective in the neutrino sector.

Anarchy and its variants, all sharing the dominance of randomness in the lepton sector, are to be confronted with models with a richer dynamical structure, some based on continuous groups [47] but in particular those based on discrete flavour groups (for reviews, see, for example, Refs. [48-50]). After the measurement of a relatively large value for θ_{13} there has been an intense work to interpret these new results along different approaches and ideas, as discussed in the talk by Smirnov [16]. Examples are suitable modifications of the minimal models [51, 52] (we discuss the Lin model of Ref. [52] in the following), modified sequential dominance models [53], larger symmetries that already at LO lead to non vanishing θ_{13} and non maximal θ_{23} [54], smaller symmetries that leave more freedom [55], models where the flavour group and a generalised CP transformation are combined in a non trivial way [56] (other approaches to discrete symmetry and CP violation are found in Refs. [57]).

Among the models with a non trivial dynamical structure those based on discrete flavour groups were motivated by the fact that the data suggest some special mixing patterns as good first approximations like Tri-Bimaximal (TB) or Golden Ratio (GR) or Bi-Maximal (BM) mixing, for example. The corresponding mixing matrices all have $\sin^2 \theta_{23} = 1/2$, $\sin^2 \theta_{13} = 0$, values that are good approximations to the data (although less so since the most recent data), and differ by the value of the solar angle $\sin^2 \theta_{12}$. The observed $\sin^2 \theta_{12}$, the best measured mixing angle, is very close, from below, to the so called Tri-Bimaximal (TB) value [58] of $\sin^2 \theta_{12} = 1/3$. Alternatively, it is also very close, from above, to the Golden Ratio (GR) value [59] $\sin^2 \theta_{12} = \frac{1}{\sqrt{5}\phi} = \frac{2}{5+\sqrt{5}} \sim 0.276$, where $\phi = (1 + \sqrt{5})/2$ is the GR (for a different connection to the GR, see Refs. [60]). On a different perspective, one has also considered models with Bi-Maximal (BM) mixing, where at leading order (LO), before diagonalization of charged leptons, $\sin^2 \theta_{12} = 1/2$, i.e. it is also maximal, and the necessary, rather large, corrective terms to θ_{12} arise from the diagonalization of the charged lepton mass matrices (a list of references can be found in Ref. [48]). Thus, if one or the other of these coincidences is taken seriously, models where TB or GR or BM mixing is naturally predicted provide a good first approximation (but these hints cannot all be relevant and it is well possible that none is). As the corresponding mixing matrices have the form of rotations with fixed special angles one is naturally led to discrete flavour groups.

In the following we will mainly refer to TB or BM mixing which are the most studied first approximations to the data. A simplest discrete symmetry for TB mixing is A_4 while BM can be obtained from S_4 . Starting with the ground breaking paper in Ref. [61], A_4 models have been widely studied (for a recent review and a list of references, see Ref. [66]). At LO the typical A_4 model (like, for example, the one discussed in in Ref. [62]) leads to exact TB mixing. In these models the starting LO approximation is completely fixed (no

chance), but the Next to LO (NLO) corrections still introduce a number of undetermined parameters, although in general much less numerous than for $U(1)_{FN}$ models. These models are therefore more predictive and in each model, one obtains relations among the departures of the three mixing angles from the LO patterns, restrictions on the CP violation phase δ_{CP} , mass sum rules among the neutrino mass eigenvalues, definite ranges for the neutrinoless beta decay effective Majorana mass and so on. Given the set of flavour symmetries and having specified the field content, the non-leading corrections to TB mixing, arising from higher dimensional effective operators, can be evaluated in a well-defined expansion. In the absence of specific dynamical tricks, in a generic model all three mixing angles receive corrections of the same order of magnitude. Since the experimentally allowed departures of θ_{12} from the TB value, $\sin^2 \theta_{12} = 1/3$, are small, numerically not larger than $\mathcal{O}(\lambda_C^2)$ where $\lambda_C = \sin \theta_C$, it follows that both θ_{13} and the deviation of θ_{23} from the maximal value are also expected to be typically of the same general size. This generic prediction of a small θ_{13} , numerically of $\mathcal{O}(\lambda_C^2)$, is at best marginal after the recent measurement of θ_{13} .

Of course, one can introduce some additional theoretical input to improve the value of θ_{13} [67]. In the case of A_4 , one particularly interesting example is provided by the Lin model [52] (see also Ref. [51]), formulated before the recent θ_{13} results. In the Lin model the A_4 symmetry breaking is arranged, by suitable additional Z_n parities, in a way that the corrections to the charged lepton and the neutrino sectors are kept separated not only at LO but also at NLO. As a consequence, in a natural way the contribution to neutrino mixing from the diagonalization of the charged leptons can be of $\mathcal{O}(\lambda_C^2)$, while those in the neutrino sector of $\mathcal{O}(\lambda_C)$. Thus, in the Lin model the NLO corrections to the solar angle θ_{12} and to the reactor angle θ_{13} are not necessarily related. In addition, in the Lin model the largest corrections do not affect θ_{12} and satisfy the relation $\sin^2 \theta_{23} = 1/2 + 1/\sqrt{2} \cos \delta_{CP} |\sin \theta_{13}|$, with δ_{CP} being the CKM-like CP violating phase of the lepton sector. Note that, for θ_{23} in the first octant, the sign of $\cos \delta_{CP}$ must be negative.

Alternatively, one can think of models where, because of a suitable symmetry, BM mixing holds in the neutrino sector at LO and the corrective terms for θ_{12} , which in this case are required to be large, arise from the diagonalization of charged lepton masses. These terms from the charged lepton sector, numerically of order $\mathcal{O}(\lambda_C)$, would then generically also affect θ_{13} and the resulting angle could well be compatible with the measured value. An explicit model of this type based on the group S_4 has been developed in Ref. [64] (see also Refs. [65]). An important feature of this particular model is that only θ_{12} and θ_{13} are corrected by terms of $\mathcal{O}(\lambda_C)$ while θ_{23} is unchanged at this order. This model is compatible with present data and clearly prefers the upper range of the present experimental result for θ_{13} . Note however that the present bounds on lepton flavour violating (LFV) reactions [68] pose severe constraints on the parameter space of the models, assuming a supersymmetric context (for a recent general analysis of LFV effects in the context of flavour models, see Ref. [69]). In particular, we refer to the recent improved MEG result [70] on the $\mu \rightarrow e\gamma$ branching ratio, $Br(\mu \rightarrow e\gamma) \leq 5.7 \times 10^{-13}$ at 90% C.L. and to other similar processes like $\tau \rightarrow (e \text{ or } \mu)\gamma$. Particularly constrained are the models with relatively large corrections from the off-diagonal terms of the charged lepton mass matrix, like the models with BM mixing at LO [67]. A way out is to push the s-partners at large enough masses but then a supersymmetric explanation of the muon (g-2) anomaly becomes less plausible [71, 72].

In conclusion, one could have imagined that neutrinos would bring a decisive boost towards the formulation of a comprehensive understanding of fermion masses and mixings. In reality it is frustrating that no real illumination was sparked on the problem of flavour. We can reproduce in many different ways the observations, in a wide range that goes from anarchy to discrete flavour symmetries but we have not yet been able to single out a unique and convincing baseline for the understanding of fermion masses and mixings. In spite of many interesting ideas and the formulation of many elegant models the mysteries of the flavour structure of the three generations of fermions have not been much unveiled.

8. – Conclusion

Bruno Pontecorvo made seminal contributions to neutrino physics. This domain of physics deals with fundamental issues still of great importance. Our knowledge of neutrino physics has been much advanced in the last 15 years and it is still vigorously studied and progress is continuously made, but many crucial problems are still open. Together with LHC physics [73] the study of neutrino and flavour processes maintains a central role in fundamental physics.

* * *

I thank the Organizers for their kind invitation and financial support. I recognize that this work has been partly supported by the COFIN program (PRIN 2008) and by the European Commission, under the networks LHCPHENONET and Invisibles

REFERENCES

- [1] See the talks by G. Mitselmakher, J. Steinberger, S. Bilenki and J. Bernabeu, these Proceedings.
- [2] G. Bellini, these Proceedings.
- [3] F. Halzen, these Proceedings.
- [4] G. Altarelli and F. Feruglio, *New J. Phys.* **6** (2004) 106, arXiv:hep-ph/0405048; R. N. Mohapatra and A. Y. Smirnov, *Ann. Rev. Nucl. Part. Sci.* **56** (2006) 569, arXiv:hep-ph/0603118; W. Grimus, *PoS P2GC* (2006) 001, arXiv:hep-ph/0612311; M. C. Gonzalez-Garcia and M. Maltoni, *Phys. Rept.* **460** (2008) 1, arXiv:0704.1800.
- [5] P. Minkowski, *Phys. Letters B* **67** (1977) 421; T. Yanagida, in *Proc. of the Workshop on Unified Theory and Baryon Number in the Universe*, KEK, March 1979; S. L. Glashow, in “Quarks and Leptons”, Cargèse, ed. M. Lévy et al., Plenum, 1980 New York, p. 707; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, Stony Brook, Sept 1979; R. N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44** (1980) 912.
- [6] Y. Wang, these Proceedings.
- [7] K. Nishikawa, these Proceedings.
- [8] G. Fogli *et. al.*, arXiv:1205.5254.
- [9] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado, T. Schwetz, arXiv:1209.3023.
- [10] D. Forero, M. Tortola, and J. Valle, arXiv:1205.4018.
- [11] C. Giunti, these Proceedings.
- [12] A. McDonald, these Proceedings.
- [13] Y. Suzuki, these Proceedings.

- [14] B. Pontecorvo, Sov. Phys. JETP **6**, 429 (1957) [Zh. Eksp. Teor. Fiz. **33**, 549 (1957)]; Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870; B. Pontecorvo, Sov. Phys. JETP **26**, 984 (1968) [Zh. Eksp. Teor. Fiz. **53**, 1717 (1968)]; V. N. Gribov and B. Pontecorvo, Phys. Lett. B **28**, 493 (1969).
- [15] B. W. Lee, S. Pakvasa, R. Shrock, and H. Sugawara, Phys. Rev. Lett. **38**, 937 (1977); B. W. Lee and R. Shrock, Phys. Rev. D **16**, 1444 (1977).
- [16] A. Y. Smirnov, these Proceedings.
- [17] N. Cabibbo, Phys. Lett. B **72**, 333 (1978).
- [18] S. M. Bilenky, J. Hosek and S. T. Petcov, Phys. Lett. B **94** (1980) 495; J. Schechter and J. W. F. Valle, Phys. Rev. D **22** (1980) 2227; M. Doi, T. Kotani, H. Nishiura, K. Okuda and E. Takasugi, Phys. Lett. B **102** (1981) 323. Mass matrices with a general dependence on ϕ , $\phi_{1,2}$ have been analyzed in: M. Frigerio and A. Y. Smirnov, Nucl. Phys. B **640** (2002) 233 and Phys. Rev. D **67** (2003) 013007.
- [19] S. Geer, these Proceedings.
- [20] C. Weinheimer, these Proceedings.
- [21] S. Hannestad, these Proceedings.
- [22] Planck Collaboration, P. A. R. Ade *et al.*, arXiv:1303.5076.
- [23] E. Fiorini, these Proceedings.
- [24] For a review see, for example,; M. Trodden, Rev. Mod. Phys. **71**, 1463 (1999).
- [25] For a review see for example: W. Buchmuller, R.D. Peccei and T. Yanagida, Ann.Rev.Nucl.Part.Sci.55, 311 ,2005. arXiv:hep-ph/0502169.
- [26] E. Giusarma *et al.*, arXiv:1102.4774; S. Joudaki *et al.*, arXiv:1208.4354.
- [27] K. Abazajian *et al.*, arXiv: 1204.5379.
- [28] C. Rubbia *et al.*, arXiv:1304.2047.
- [29] MiniBooNE Collaboration, A. A. Aguilar-Arevalo *et al.*, arXiv:1207.4809; arXiv:1303.2588.
- [30] ICARUS Collaboration, M. Antonello *et al.*, arXiv:1209.0122.
- [31] G. Mention *et al.*, arXiv:1101.2755.
- [32] P. Huber, arXiv:1106.0687.
- [33] C. Giunti and M. Laveder, arXiv:1006.3244.
- [34] J. Kopp, P. A.N. Machado, M. Maltoni and T. Schwetz, arXiv:1303.3011.
- [35] C. Giunti, M. Laveder, Y.F. Li, H.W. Long, arXiv:1308.5288; C. Giunti, arXiv:1311.1335.
- [36] J. M. Conrad *et al*, arXiv:1207.4765.
- [37] ICARUS and NESSIE Coll., M. Antonello *et al.*, arXiv:1203.3432.
- [38] M. Archidiacono *et al.*, arXiv:1302.6720.
- [39] A. Palazzo, arXiv:1302.1102.
- [40] A. Mirizzi *et al.*, arXiv:1303.5368.
- [41] L. J. Hall, H. Murayama, and N. Weiner, Phys. Rev. Lett. **84** (2000) 2572, arXiv:hep-ph/9911341.
- [42] A. de Gouvea and H. Murayama, Phys. Lett. **B573** (2003) 94, arXiv:hep-ph/0301050.
- [43] A. de Gouvea and H. Murayama, arXiv:1204.1249.
- [44] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. **B147** (1979) 277.
- [45] G. Altarelli, F. Feruglio, I. Masina and L. Merlo, arXiv:1207.0587.
- [46] W. Buchmuller, V. Domcke, and K. Schmitz, , JHEP **03** (2012) 008, arXiv:1111.3872.
- [47] See, for example, I. d. M. Varzielas and G. G. Ross, arXiv:1203.6636; R. Alonso *et al.*, JHEP **06** (2011) 037, arXiv:1103.5461; R. Alonso, M. Gavela, D. Hernandez, and L. Merlo, Phys.Lett. B **715** (2012) 194, arXiv:1206.3167; G. Blankenburg, G. Isidori and J. Jones-Perez, arXiv:1204.0688; R. Alonso, M. B. Gavela, G. Isidori and L. Maiani, arXiv:1306.5927.
- [48] G. Altarelli and F. Feruglio, , Rev. Mod. Phys. **82** (2010) 2701, arXiv:1002.0211.
- [49] H. Ishimori *et al.*, Prog. Theor. Phys. Suppl. **183** (2010) 1, arXiv:1003.3552; S. King and C. Luhn, arXiv:1301.1340.
- [50] W. Grimus and P. O. Ludl, J. Phys. A **45** (2012) 233001, arXiv:1110.6376.

- [51] E. Ma and D. Wegman, Phys. Rev. Lett. 107 (2011) 061803, arXiv:1106.4269; S. F. King and C. Luhn, JHEP 09 (2011) 042, arXiv:1107.5332; F. Bazzocchi, arXiv:1108.2497; I. de Medeiros Varzielas and L. Merlo, JHEP 02 (2011) 062, arXiv:1011.6662; W. Rodejohann and H. Zhang, arXiv:1207.1225; F. Bazzocchi and L. Merlo, arXiv:1205.5135.
- [52] Y. Lin, Nucl. Phys. **B824** (2010) 95, arXiv:0905.3534.
- [53] See, for example, S.F. King, arXiv:1311.3295.
- [54] R. de Adelhart Toorop, F. Feruglio, and C. Hagedorn, Phys. Lett. B703 (2011) 447, arXiv:1107.3486; Nucl. Phys. B858 (2012) 437, arXiv:1112.1340; S. F. King, C. Luhn, A. J. Stuart, arXiv:1207.5741; C. Hagedorn and D. Meloni, Nucl. Phys. B 862 (2012) 691, arXiv:1204.0715; T. Araki et al, arXiv:1309.4217; M. Holthausen and K. S. Lim, Phys. Rev. D 88 (2013) 033018, arXiv:1306.4356; S. F. King, T. Neder and A. J. Stuart, Phys. Lett. B 726 (2013) 312, arXiv:1305.3200.
- [55] S. -F. Ge, D. A. Dicus and W. W. Repko, Phys. Lett. B 702 (2011) 220, arXiv:1104.0602; Phys. Rev. Lett. 108 (2012) 041801, arXiv:1108.0964; D. Hernandez and A. Y. Smirnov, Phys. Rev. D 86 (2012) 053014, arXiv:1204.0445.
- [56] G. Ecker, W. Grimus and H. Neufeld, J. Phys. A 20 (1987) L807; Int. J. Mod. Phys. A 3 (1988) 603; W. Grimus and L. Lavoura, Phys. Lett. B 579 (2004) 113, arXiv:hep-ph/0305309; R. Krishnan, P. F. Harrison and W. G. Scott, arXiv:1211.2000; R. N. Mohapatra and C. C. Nishi, Phys. Rev. D 86 (2012) 073007, arXiv:1208.2875; M. Holthausen, M. Lindner and M. A. Schmidt, arXiv:1211.6953; W. Grimus and M. N. Rebelo, Phys. Rept. 281 (1997) 239, arXiv:hep-ph/9506272; F. Feruglio, C. Hagedorn and R. Ziegler, arXiv:1211.5560; arXiv:1303.7178.
- [57] G. C. Branco, J. M. Gerard and W. Grimus, Phys. Lett. B 136 (1984) 383; I. de Medeiros Varzielas and D. Emmanuel-Costa, Phys. Rev. D 84 (2011) 117901; arXiv:1106.5477; I. de Medeiros Varzielas, D. Emmanuel-Costa and P. Leser, Phys. Lett. B 716 (2012) 193; arXiv:1204.3633; I. de Medeiros Varzielas, JHEP 1208 (2012) 055; arXiv:1205.3780; G. Bhattacharyya, I. de Medeiros Varzielas and P. Leser, Phys. Rev. Lett. 109 (2012) 241603; arXiv:1210.0545; K. S. Babu and J. Kubo, Phys. Rev. D 71 (2005) 056006; arXiv:hep-ph/0411226; K. S. Babu, K. Kawashima and J. Kubo, Phys. Rev. D 83 (2011) 095008, arXiv:1103.1664; M. -C. Chen and K. T. Mahanthappa, Phys. Lett. B 681 (2009) 444; arXiv:0904.1721; A. Meroni, S. T. Petcov and M. Spinrath, Phys. Rev. D 86 (2012) 113003, arXiv:1205.5241.
- [58] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B530** (2002) 167, arXiv:hep-ph/0202074; P. F. Harrison and W. G. Scott, , Phys. Lett. **B535** (2002) 163, arXiv:hep-ph/0203209; Z.-z. Xing, Phys. Lett. **B533** (2002) 85, arXiv:hep-ph/0204049; P. F. Harrison and W. G. Scott, Phys. Lett. **B547** (2002) 219, arXiv:hep-ph/0210197; Phys. Lett. **B557**(2003) 76, arXiv:hep-ph/0302025.
- [59] Y. Kajiyama, M. Raidal, and A. Strumia, Phys. Rev. **D76** (2007) 117301, arXiv:0705.4559; L. L. Everett and A. J. Stuart, Phys. Rev. **D79** (2009) 085005, arXiv:0812.1057; G.-J. Ding, L. L. Everett, and A. J. Stuart, Nucl. Phys. **B857** (2012) 219, arXiv:1110.1688; F. Feruglio and A. Paris, JHEP **03** (2011) 101, arXiv:1101.0393.
- [60] W. Rodejohann, Phys. Lett. **B671** (2009) 267, arXiv:0810.5239; A. Adulpravitchai, A. Blum, and W. Rodejohann, New J. Phys. **11** (2009) 063026, arXiv:0903.0531.
- [61] E. Ma and G. Rajasekaran, Phys. Rev. **D64**(2001) 113012, arXiv:hep-ph/0106291.
- [62] G. Altarelli and F. Feruglio, Nucl. Phys. **B741** (2006) 215, arXiv:hep-ph/0512103.
- [63] G. Altarelli, F. Feruglio, and I. Masina, Nucl. Phys. **B689** (2004) 157, arXiv:hep-ph/0402155.
- [64] G. Altarelli, F. Feruglio, and L. Merlo, JHEP **05** (2009) 020, arXiv:0903.1940.
- [65] R. de Adelhart Toorop, F. Bazzocchi, and L. Merlo, JHEP **08** (2010) 001, arXiv:1003.4502; K. M. Patel, Phys. Lett. **B695** (2011) 225, arXiv:1008.5061; D. Meloni, JHEP **10** (2011) 010, arXiv:1107.0221.
- [66] G. Altarelli, F. Feruglio, and L. Merlo, arXiv:1205.5133.
- [67] G. Altarelli, F. Feruglio, L. Merlo and E. Stamou, JHEP 1208 (2012) 021, arXiv:1205.4670.
- [68] A. Masiero, these Proceedings.

- [69] L. Calibbi, Z. Lalak, S. Pokorski, and R. Ziegler, arXiv:1204.1275.
- [70] J. Adam et al, [MEG Collaboration], arXiv:1303.0754.
- [71] Muon g-2 Collab., G. W. Bennett et al, Phys. Rev. D73, 072003 (2006); B. L. Roberts, Chin. Phys. C34, 741 (2010).
- [72] A. Hoecker and W. Marciano, The Muon Anomalous Magnetic Moment, in Particle Data Group, J. Beringer et al, Phys. Rev. D86(2012) 010001.
- [73] See the talks by R. Barbieri and S. Bertolucci, these Proceedings.