

## The MUonE experiment: A measurement of the hadronic contribution to the muon $g - 2$ via $\mu$ - $e$ elastic scattering

R. N. PILATO<sup>(1)</sup><sup>(2)</sup> on behalf of the MUONE COLLABORATION

<sup>(1)</sup> *INFN, Sezione di Pisa - Pisa, Italy*

<sup>(2)</sup> *Dipartimento di Fisica, Università di Pisa - Pisa, Italy*

received 8 June 2020

**Summary.** — We present the MUonE experimental proposal, which aims to determine the leading-order hadronic contribution to the muon  $g - 2$  using a novel approach, based on the measurement of the hadronic contribution to the running of the electromagnetic coupling constant in the space-like region.

### 1. – Introduction

The measurement of the muon magnetic anomaly,  $a_\mu = (g_\mu - 2)/2$ , presently exhibits a  $\sim 3.7\sigma$  discrepancy from the Standard Model prediction [1], representing a possible hint of new physics. On the experimental side,  $a_\mu$  will be measured in the next years at the remarkable accuracy of  $\sim 0.14$  ppm by two new experiments at Fermilab and J-PARC, improving by a factor of 4 the precision of the most recent result. An improvement is also required on the theoretical prediction, as its uncertainty can become the main limitation for a test of the Standard Model. The accuracy on the Standard Model calculation is limited by the evaluation of the leading-order hadronic contribution  $a_\mu^{HLO}$ , which cannot be computed perturbatively at low energies. For this reason,  $a_\mu^{HLO}$  is traditionally determined by means of a dispersion integral on the annihilation cross section  $e^+e^- \rightarrow$  hadrons, which is densely populated by resonances and influenced by flavour threshold effects. These aspects limit the final precision achievable by this method. Nevertheless, the calculation of  $a_\mu^{HLO}$  has reached the accuracy of  $\sim 0.4\%$ . In order to consolidate the theoretical prediction, it is important to crosscheck this calculation in an independent way.

### 2. – The MUonE experiment

The MUonE experiment has been recently proposed [2], with the aim to measure  $a_\mu^{HLO}$  using a completely independent approach. It is based on the measurement of the




Fig. 1. – Sketch of a single station (image not to scale).

hadronic contribution to the running of the electromagnetic coupling constant ( $\Delta\alpha_{had}$ ) in the space-like region, by means of  $\mu^\pm e^- \rightarrow \mu^\pm e^-$  elastic scattering. The measurement of the shape of the differential cross section provides direct sensitivity to  $\Delta\alpha_{had}$ , and it is carried out by scattering a 150 GeV muon beam on a beryllium target. The M2 beam available at CERN provides muons with the proper energy and an average intensity of  $1.3 \cdot 10^7 \mu/s$ . It allows to collect an integrated luminosity of  $1.5 \times 10^7 \text{ nb}^{-1}$  in 3 years of data taking, corresponding to a statistical uncertainty of 0.3% on  $a_\mu^{HLO}$ . This makes the measurement of MUonE competitive with the dispersive approach.

**2.1. Experimental apparatus.** – The experimental apparatus consists of a repetition of 40 identical stations. A sketch of a single station is shown in fig. 1. It is made up of a 15 mm thick beryllium target, followed by a tracking system with a lever arm of  $\sim 1$  m, which is used to measure the scattering angles with high precision. The tracking system is composed by 3 pairs of silicon strip sensors. In particular, the sensors foreseen for the CMS HL-LHC Outer Tracker in the so-called 2S configuration have been chosen. A sensor is made up of two layers reading the same coordinate. Each layer is  $320 \mu\text{m}$  thick, with a squared area of  $10 \times 10 \text{ cm}^2$  and a pitch of  $90 \mu\text{m}$ , which allows to obtain an angular resolution of  $\sim 20 \mu\text{rad}$ . The apparatus is also equipped with an electromagnetic calorimeter, placed downstream all the stations. Its main role is to provide  $e/\mu$  particle identification. The final optimization of the calorimeter is still under study. Two options are currently considered:  $\text{PbWO}_4$  and  $\text{PbF}_2$  crystals. A surface of  $\sim 1 \times 1 \text{ m}^2$  will allow to achieve a full acceptance for electrons in the angular region of interest (scattering angles  $\lesssim 10 \text{ mrad}$ ).

**2.2. Systematic uncertainties.** – The main challenge of the MUonE experiment is to reach a systematic uncertainty of the same order as the statistical one. For this purpose, the differential cross section must be measured with a systematic uncertainty  $\lesssim 10 \text{ ppm}$ . Systematic uncertainties arise both from experimental and theoretical aspects, such as: bad reconstruction of the elastic events, limited control of the experimental conditions, missing contributions in the computation of the theoretical cross section.

**2.3. Future plans.** – A Letter of Intent has been submitted in June 2019 to CERN SPSC [3]. Studies on detector optimization, simulations and theory improvements will continue in 2020. The detector construction is expected during CERN LS2 and the plan is to have a pilot run of 3 weeks in 2021. A run with full statistics is envisaged in 2022–24.

## REFERENCES

- [1] KESHAVARZI A., NOMURA D. and TEUBNER T., *Phys. Rev. D*, **97** (2018) 114025.
- [2] ABBIENDI G. *et al.*, *Eur. Phys. J. C*, **77** (2017) 139.
- [3] THE MUonE COLLABORATION, *Letter of Intent: the MUonE project*, CERN-SPSC-2019-026 (2019).